

A service of the U.S. National Library of Medicine

NIH National Institutes of Health

Home About MedlinePlus Site Map FAQs Contact Us

Search MedlinePlus

GO

**Health Topics** 

Drugs & Supplements

Videos & Cool Tools

ESPAÑOL

# Hernia

f 💆 🛨

A hernia is a sac formed by the lining of the abdominal cavity (peritoneum). The sac comes through a hole or weak area in the strong layer of the belly wall that surrounds the muscle. This layer is called the fascia.

Which type of hernia you have depends on where it is:

- Femoral hernia is a bulge in the upper thigh, just below the groin. This type is more common in women than men.
- Hiatal hernia occurs in the upper part of the stomach. Part of the upper stomach pushes into the chest.
- Incisional hernia can occur through a scar if you have had abdominal surgery in the past.
- <u>Umbilical hernia</u> is a bulge around the belly button. It occurs when the muscle around the belly button
  does not close completely.
- Inguinal hernia is a bulge in the groin. It is more common in men. It may go all the way down into the scrotum.

## Causes

Usually, there is no clear cause of a hernia. Sometimes, hernias occur with heavy lifting, straining while using the toilet, or any activity that raises the pressure inside the belly. Hernias may be present at birth, but the bulge may not be noticeable until later in life. Some persons have a family history of hernias.

Babies and children can get hernias. It happens when there is weakness in the belly wall. About 5 out of 100 children have inguinal hernias. They are more common in boys. Some children do not have symptoms until they are adults.

Any activity or medical problem that increases pressure on the tissue in the belly wall and muscles may lead to a hernia, including:

- · Chronic constipation and pushing hard (straining) to have a bowel movement
- · Chronic cough
- Cystic fibrosis
- Enlarged prostate, straining to urinate
- · Extra weight
- Fluid in the abdomen (ascites)
- · Heavy lifting
- Peritoneal dialysis
- Poor nutrition
- Smoking
- Overexertion
- Undescended testicles

## **Symptoms**

There are usually no symptoms. Some people have discomfort or pain. The discomfort may be worse when standing, straining, or lifting heavy objects. In time, the most common complaint is a bump that is sore and growing.

When a hernia gets bigger, it may get stuck inside the hole and lose its blood supply. This is called strangulation. Symptoms include nausea and vomiting and not being able to pass gas or have bowel movements. When this occurs, surgery is needed right away.

## **Exams and Tests**

The health care provider can usually see or feel a hernia when you are examined. You may be asked to

# MedlinePlus Topics

**Hernia** 

# **Images**





Inguinal hernia

Inguinal hernia repair - Series

## Read More

Femoral hernia
Femoral hernia repair
Inguinal hernia repair
Lump in the abdomen
Umbilical hernia repair

# Patient Instructions

Inguinal hernia - discharge

cough, bend, push, or lift. The hernia may get bigger when you do this.

The hernia (bulge) may not be easily seen in infants and children, except when the child is crying or coughing.

<u>Ultrasound</u> or <u>CT scan</u> may be done to look for a hernia.

If there is a blockage in the bowel, an x-ray of the abdomen will likely be done.

#### Treatment

Surgery is the only treatment that can permanently fix a hernia. Surgery may be more risky for patients with serious medical problems.

Surgery secures the weakened abdominal wall tissue (fascia) and closes any holes. Most hernias are closed with cloth patches to plug the hole.

An umbilical hernia that does not heal on its own by the time a child is 5 years old will likely be repaired.

## Outlook (Prognosis)

The outcome for most hernias is usually good with treatment. It is rare for a hernia to come back. Incisional hernias are more likely to return.

#### Possible Complications

In rare cases, inguinal hernia repair can damage structures involved in the function of a man's testicles.

Another risk of hernia surgery is nerve damage, which can lead to numbness in the groin area.

If a part of the bowel was trapped or strangulated before surgery, bowel perforation or dead bowel may result.

## When to Contact a Medical Professional

Call your health care provider right away if:

- You have a painful hernia and the contents cannot be pushed back into the abdomen using gentle
  pressure
- · You develop nausea, vomiting, or a fever along with a painful hernia
- You have a hernia that becomes red, purple, dark, or discolored

Call your health care provider if:

- · You have groin pain, swelling, or a bulge
- You have a bulge or swelling in the groin or belly button, or that is associated with a previous surgical
  cut.

## Prevention

- Use proper lifting techniques.
- Lose weight if you are overweight.
- Relieve or avoid constipation by eating plenty of fiber, drinking lots of fluid, going to the bathroom as soon as you have the urge, and exercising regularly.
- Men should see their health care provider if they strain with urination. This may be a symptom of an enlarged prostate.

# Alternative Names

Hernia - inguinal; Inguinal hernia; Direct and indirect hernia; Rupture; Strangulation; Incarceration

### References

Jeyarajah DR, Harford WV Jr. Abdominal hernias and gastric volvulus. In: Feldman M, Friedman LS, Brandt LJ, eds. Sleisenger and Fordtran's Gastrointestinal and Liver Disease

Pathophysiology/Diagnosis/Management. 9th ed. Philadelphia, Pa: Elsevier Saunders; 2010:chap 24.

Malangoni MA, Rosen MJ. Hernias. In: Townsend CM, Beauchamp RD, Evers BM, Mattox KL, eds. *Sabiston Textbook of Surgery*. 19th ed. Philadelphia, Pa: Elsevier Saunders; 2012:chap 46.

Update Date: 11/15/2013

Updated by: Debra G. Wechter, MD, FACS, General Surgery practice specializing in breast cancer, Virginia Mason Medical Center, Seattle, Washington. Also reviewed by David Zieve, MD, MHA, Bethanne Black, and the A.D.A.M. Editorial team.

#### Browse the Encyclopedia



A.D.A.M., Inc. is accredited by URAC, also known as the American Accreditation HealthCare Commission (www.urac.org). URAC's <u>accreditation program</u> is an independent audit to verify that A.D.A.M. follows rigorous standards of quality and accountability. A.D.A.M. is among the first to achieve this important distinction for online health information and services. Learn more about A.D.A.M.'s <u>editorial policy</u>, <u>editorial process</u> and <u>privacy policy</u>. A.D.A.M. is also a founding member of Hi-Ethics and subscribes to the principles of the Health on the Net Foundation (www.hon.ch).

The information provided herein should not be used during any medical emergency or for the diagnosis or treatment of any medical condition. A licensed physician should be consulted for diagnosis and treatment of any and all medical conditions. Call 911 for all medical emergencies. Links to other sites are provided for information only -- they do not constitute endorsements of those other sites. Copyright 1997-2014, A.D.A.M., Inc. Duplication for commercial use must be authorized in writing by ADAM Health Solutions.



<u>Disclaimers</u> <u>Copyright</u> <u>Privacy</u> <u>Accessibility</u> <u>Quality Guidelines</u> <u>Viewers & Players</u>

U.S. National Library of Medicine 8600 Rockville Pike, Bethesda, MD 20894 U.S. Department of Health and Human Services National Institutes of Health

Page last updated: 16 May 2014